



General License Class

Chapter 8 Propagation



The Ionosphere

- Regions
 - Ionosphere.
 - A region of the atmosphere extending from 30 miles to 300 miles above the surface of the earth.
 - Solar radiation causes atoms in the ionosphere to become ionized.




The Ionosphere

- Regions
 - Ionosphere.
 - The ionosphere organizes itself into regions or “layers”.
 - Varies with strength of ionization.
 - D-region disappears at night.
 - F-region splits into F₁ & F₂ regions during the day.



The Ionosphere

- Regions
 - D-Layer.
 - 30-60 miles altitude.
 - Rapidly disappears at sunset.
 - Rapidly re-forms at sunrise.
 - Absorbs long wavelength radio waves.
 - 160m, 80m, & 40m.






The Ionosphere

- Regions

- E-Layer.

- 60-70 miles altitude.
 - One hop up to 1200 miles.
 - Similar to D-layer.
 - Lasts longer into the night.
 - Less absorption.
 - Auroral propagation.
 - Sporadic-E skip.
 - 10m, 6m, & 2m.




The Ionosphere

- Regions

- F-Layer.


- 100-300 miles altitude.
 - One-hop up to 2500 miles.
 - Can remain ionized all night.
 - Splits into F1-layer & F2-layer during the day.
 - F1-layer = 100-140 miles.
 - F2-layer = 200-300 miles.
 - Long-range HF propagation.





The Ionosphere

- Reflection & Absorption
 - Radio waves are refracted (bent) in the ionosphere.



The Ionosphere

- Reflection & Absorption
 - Radio waves are refracted (bent) in the ionosphere.
 - The stronger the ionization level, the more the waves will be bent.
 - The shorter the wavelength (higher frequency), the less the waves will be bent.
 - VHF & UHF are only slightly bent & almost never enough to return to earth.



The Ionosphere

- Reflection & Absorption
 - Radio waves are refracted (bent) in the ionosphere.
 - Critical angle.
 - Maximum angle at which radio waves are bent enough to return to earth.
 - Critical angle decreases with increasing frequency.
 - One reason why a low angle of radiation is important for working DX.



The Ionosphere

- Reflection & Absorption
 - Radio waves are refracted (bent) in the ionosphere.
 - Critical frequency.
 - Highest frequency at which radio waves sent straight up are bent enough to return to earth.
 - Important for NVIS operation.



The Ionosphere

- Reflection & Absorption
 - Absorption.
 - Atmosphere is denser at lower altitudes, causing part of the RF energy to be absorbed.
 - The longer the wavelength (lower frequency), the higher the amount of absorption.




The Ionosphere

- Reflection & Absorption
 - Absorption.
 - D-region.
 - Almost no refraction (bending) of radio waves.
 - Almost completely absorbs radio waves below 10 MHz.
 - E-region.
 - More refraction than D-region.
 - Less absorption than D-region.



The Ionosphere

- Reflection & Absorption



The Ionosphere

- Sky-Wave & Ground-Wave Propagation.
 - Sky-Wave.
 - Refracting radio waves back to earth using the ionosphere (a.k.a. – skip).
 - Each trip from earth to ionosphere & back to earth is called a "hop".
 - Multiple hops are common.



The Ionosphere

- Sky-Wave & Ground-Wave Propagation.
 - Sky-Wave.
 - Maximum distance of a single hop depends on altitude of the region where refraction takes place.
 - E-region = Single hop can be up to 1200 miles.
 - F-region = Single hop can be up to 2500 miles.



The Ionosphere

- Sky-Wave & Ground-Wave Propagation.
 - Sky-Wave.
 - Hops considerably less than the maximum distance is called short skip.
 - Higher critical angle
 - Short skip is a good indicator that skip is possible on a higher frequency band.
 - Fluttery sound result of irregularities in ionization region causing multiple paths.



The Ionosphere

- Sky-Wave & Ground-Wave Propagation.
 - Ground-Wave.
 - Radio waves can follow along the surface of the earth.
 - Primarily vertically polarized.
 - Losses in earth cause rapid decrease of signal strength for increasing distance.
 - Higher frequency → higher loss.




The Ionosphere

- Sky-Wave & Ground-Wave Propagation.
 - Skip distance.
 - Distance from transmitter where radio wave first returns to earth.
 - Skip zone.
 - Too close for sky-wave propagation, but too far for ground-wave propagation.



The Ionosphere

- Sky-Wave & Ground-Wave Propagation.



The Ionosphere

- Long Path & Short Path
 - Short path.
 - Direct route between stations
 - Shortest distance.
 - Long path.
 - 180° from short path.
 - Longest distance.



The Ionosphere

- Long Path & Short Path
 - Conditions may not support short path, but long path may be possible.
 - Echo indicates both short & long paths are open.



The Ionosphere

- Long Path & Short Path





G2D06 -- How is a directional antenna pointed when making a “long-path” contact with another station?

- A. Toward the rising Sun
- B. Along the gray line
- ➔ C. 180 degrees from its short-path heading
- D. Toward the north



G3B01 -- How might a sky-wave signal sound if it arrives at your receiver by both short path and long path propagation?

- A. Periodic fading approximately every 10 seconds
- B. Signal strength increased by 3 dB
- C. The signal might be cancelled causing severe attenuation
- ➔ D. A well-defined echo might be heard



G3B02 -- Which of the following is a good indicator of the possibility of sky-wave propagation on the 6 meter band?

- A. Short skip sky-wave propagation on the 10 meter band
- B. Long skip sky-wave propagation on the 10 meter band
- C. Severe attenuation of signals on the 10 meter band
- D. Long delayed echoes on the 10 meter band



G3B09 -- What is the approximate maximum distance along the Earth's surface that is normally covered in one hop using the F2 region?

- A. 180 miles
- B. 1,200 miles
- C. 2,500 miles
- D. 12,000 miles



G3B10 -- What is the approximate maximum distance along the Earth's surface that is normally covered in one hop using the E region?

- A. 180 miles
- B. 1,200 miles
- C. 2,500 miles
- D. 12,000 miles



G3C01 -- Which of the following ionospheric layers is closest to the surface of the Earth?

- A. The D layer
- B. The E layer
- C. The F1 layer
- D. The F2 layer



G3C02 -- Where on the Earth do ionospheric layers reach their maximum height?

- ➔ A. Where the Sun is overhead
- B. Where the Sun is on the opposite side of the Earth
- C. Where the Sun is rising
- D. Where the Sun has just set



G3C03 -- Why is the F2 region mainly responsible for the longest distance radio wave propagation?

- A. Because it is the densest ionospheric layer
- B. Because it does not absorb radio waves as much as other ionospheric regions
- ➔ C. Because it is the highest ionospheric region
- D. All of these choices are correct



G3C04 -- What does the term “critical angle” mean as used in radio wave propagation?

- A. The long path azimuth of a distant station
- B. The short path azimuth of a distant station
- C. The lowest takeoff angle that will return a radio wave to the Earth under specific ionospheric conditions
- ➔ D. The highest takeoff angle that will return a radio wave to the Earth under specific ionospheric conditions



G3C05 -- Why is long distance communication on the 40, 60, 80 and 160 meter bands more difficult during the day?

- A. The F layer absorbs signals at these frequencies during daylight hours
- B. The F layer is unstable during daylight hours
- ➔ C. The D layer absorbs signals at these frequencies during daylight hours
- D. The E layer is unstable during daylight hours




G3C12 -- Which ionospheric layer is the most absorbent of long skip signals during daylight hours on frequencies below 10 MHz?

- A. The F2 layer
- B. The F1 layer
- C. The E layer
- ➔ D. The D layer



The Sun


- Sunspots & Cycles
 - Sunspots.
 - Areas of intense magnetic activity on the surface (photosphere) of the sun.






The Sun

- Sunspots & Cycles
 - Sunspots.
 - Up to 50,000 miles in diameter.
 - Emit UV radiation which ionizes earth's atmosphere.
 - Earliest observation dates from 354 BC.



The Sun


- Sunspots & Cycles
 - Sunspots.
 - Cooler in temperature (4,900°F to 7,600°F) than surrounding surface (10,000°F) so they appear darker.





The Sun

- Sunspots & Cycles
 - Sunspots.
 - Emit UV radiation which ionizes earth's atmosphere.




The Sun

- Sunspots & Cycles
 - Sunspots.
 - Life span of less than a day to a few weeks.
 - Stationary on sun's surface.
 - Appear to move because of sun's rotation.
 - Sunspots rotate back into view every 28 days.




The Sun

- Sunspots & Cycles
 - Solar Cycles.
 - Number of sunspots varies in 11-year cycles.



The Sun

- Sunspots & Cycles
 - Solar Cycles.
 - At beginning of cycle, sunspots appear at mid latitudes & appear closer to equator as cycles progresses.





The Sun

- Sunspots & Cycles
 - Solar Cycles.
 - At peak of solar cycle, ionization level can be high enough that 10m stays open all night.
 - At minimum of solar cycle, bands above 20m may not be open at all.



The Sun

- Sunspots & Cycles
 - Solar Cycles.
 - Strong seasonal & daily variations in propagation.
 - Seasonal variations due to different levels of ionization between summer & winter.
 - Seasonal variations on lower bands due to lower atmospheric noise during winter months.
 - Daily variations due to different levels of ionization between day & night.



The Sun

- Measuring Solar Activity
 - Sunspot number (SSN).
 - $SSN = 10 \times \text{Nr of groups} + \text{Nr of sunspots}$.
 - Average of observations from many different locations.
 - Solar flux index (SFI).
 - Measure of 10.7 cm (2.8 GHz) solar radiation.
 - Indicator of UV radiation.
 - Minimum value = 65, no maximum.



The Sun

- Measuring Solar Activity
 - K-index (K_p).
 - Measure of short-term stability of earth's magnetic field.
 - Minimum value = 0
 - Maximum value = 9.
 - Updated every 3 hours.
 - Higher values → Poorer HF propagation.

K-Index	Meaning
0	Inactive
1	Very quiet
2	Quiet
3	Unsettled
4	Active
5	Minor storm
6	Major storm
7	Severe storm
8	Very severe storm
9	Extremely severe storm



The Sun

- Measuring Solar Activity

- A-index (A_p).

- Measure of long-term stability of earth's magnetic field.
 - Minimum value = 0, maximum value = 400.
 - Calculated from previous 8 K-index values.
 - Higher values → Poorer HF propagation.

A-Index	Meaning
0-7	Quiet
8-15	Unsettled
16-29	Active
30-49	Minor storm
50-99	Major storm
100-400	Severe storm



The Sun

- Assessing Propagation

- Maximum useable frequency (MUF).

- Highest frequency that will allow communications between 2 points.
 - Radio waves on frequencies below the MUF will be refracted back to earth.
 - Radio waves on frequencies above the MUF will be lost into space.
 - Use a frequency just below the MUF for the best results.



The Sun

- Assessing Propagation
 - Lowest useable frequency (LUF).
 - Lowest frequency that will allow communications between 2 points.
 - Radio waves on frequencies below the LUF will be absorbed by the D-region.
 - If the MUF drops below the LUF, then sky-wave communications are not possible between those 2 points.



The Sun

- Assessing Propagation
 - International beacons.
 - Transmitters placed at 18 locations around the world.
 - Sponsored by the NCDXF & the IARU.
 - 14.100 MHz, 18.110 MHz, 21.150 MHz, 24.930 MHz, & 28.200 MHz.
 - Sends callsign (22 wpm) followed by 4 1-second dashes.
 - Callsign & 1st dash = 100 Watts.
 - 2nd dash = 10 Watts.
 - 3rd dash = 1 Watt.
 - 4th dash = 0.1 Watt.




The Sun

- Assessing Propagation
 - Software.
 - VOACAP
 - Voice of America Coverage Analysis Program
 - Online predictions.
 - <http://www.voacap.com/prediction.html>
 - <http://www.voacap.com/coverage.html>
- **Ignore the predictions – just listen!**




The Sun

- Solar Disturbances
 - Solar flare.
 - A large eruption of energy & particles from surface of the sun.
 - Caused by disruptions of the sun's magnetic field.
 - Takes about 8 minutes for energy to reach earth.



The Sun


- Solar Disturbances
 - Coronal hole.
 - A weak area in the corona through which plasma can escape the sun's magnetic field & stream through space at high velocity.





The Sun

- Solar Disturbances
 - Coronal mass ejection (CME).
 - Ejection of a large amount of material from the corona.
 - Narrow beam or wide area.
 - Often associated with a large solar flare.
 - Takes about 20-40 hours for particles to reach earth.





The Sun

- Solar Disturbances
 - Sudden ionospheric disturbance (SID).
 - UV-rays & X-rays from a solar flare travel to earth at the speed of light (186,000 mi/sec).
 - Reach earth in about 8 minutes.
 - Greatly increases ionization level of D-region.
 - Lower frequencies more greatly affected.
 - Can last from a few seconds to several hours.
 - Only affects sunlit side of earth.



The Sun

- Solar Disturbances
 - Geomagnetic disturbances.
 - CME's greatly increase strength of solar wind.
 - A continuous stream of charged particles.
 - Reaches earth in about 20-40 hours.
 - Particles become trapped in the magnetosphere near both poles increasing ionization of E-region & creating a geomagnetic storm.



The Sun

- Solar Disturbances
 - Geomagnetic disturbances.
 - High-latitude HF propagation greatly decreased.
 - Can last several hours to a few days.
 - Auroral activity greatly increased.
 - Reflection possible on 15m & up.
 - Strongest on 6m & 2m.
 - Signals modulated with hiss or buzz.
 - CW.



The Sun

- Solar Disturbances





Break



G3A01 -- What is the significance of the sunspot number with regard to HF propagation?

- A. Higher sunspot numbers generally indicate a greater probability of good propagation at higher frequencies
- B. Lower sunspot numbers generally indicate greater probability of sporadic E propagation
- C. A zero sunspot number indicate radio propagation is not possible on any band
- D. All of these choices are correct.



G3A02 -- What effect does a Sudden Ionospheric Disturbance have on the daytime ionospheric propagation of HF radio waves?

- A. It enhances propagation on all HF frequencies
- ➔ B. It disrupts signals on lower frequencies more than those on higher frequencies
- C. It disrupts communications via satellite more than direct communications
- D. None, because only areas on the night side of the Earth are affected



G3A03 -- Approximately how long does it take the increased ultraviolet and X-ray radiation from solar flares to affect radio-wave propagation on the Earth?

- A. 28 days
- B. 1 to 2 hours
- ➔ C. 8 minutes
- D. 20 to 40 hours



G3A04 -- Which of the following amateur radio HF frequencies are least reliable for long distance communications during periods of low solar activity?

- A. 80 meters and 160 meters
- B. 60 meters and 40 meters
- C. 30 meters and 20 meters
- ➔ D. 15 meters, 12 meters and 10 meters



G3A05 -- What is the solar-flux index?

- A. A measure of the highest frequency that is useful for ionospheric propagation between two points on the Earth
- B. A count of sunspots which is adjusted for solar emissions
- C. Another name for the American sunspot number
- ➔ D. A measure of solar radiation at 10.7 centimeters wavelength



G3A06 -- What is a geomagnetic storm?

- A. A sudden drop in the solar-flux index
- B. A thunderstorm which affects radio propagation
- C. Ripples in the ionosphere
- D. A temporary disturbance in the Earth's magnetosphere



G3A07 -- At what point in the solar cycle does the 20 meter band usually support worldwide propagation during daylight hours?

- A. At the summer solstice
- B. Only at the maximum point of the solar cycle
- C. Only at the minimum point of the solar cycle
- D. At any point in the solar cycle



G3A08 -- Which of the following effects can a geomagnetic storm have on radio-wave propagation?

- A. Improved high-latitude HF propagation
- ➔ B. Degraded high-latitude HF propagation
- C. Improved ground-wave propagation
- D. Improved chances of UHF ducting



G3A09 -- What effect does a high sunspot number have on radio communications?

- A. High-frequency radio signals become weak and distorted
- B. Frequencies above 300 MHz become usable for long-distance communication
- ➔ C. Long-distance communication in the upper HF and lower VHF range is enhanced
- D. Microwave communications become unstable



G3A10 -- What causes HF propagation conditions to vary periodically in a 28-day cycle?

- A. Long term oscillations in the upper atmosphere
- B. Cyclic variation in the Earth's radiation belts
- C. The Sun's rotation on its axis
- D. The position of the Moon in its orbit



G3A11 -- Approximately how long is the typical sunspot cycle?

- A. 8 minutes
- B. 40 hours
- C. 28 days
- D. 11 years



G3A12 -- What does the K-index indicate?

- A. The relative position of sunspots on the surface of the Sun
- ➔ B. The short term stability of the Earth's magnetic field
- C. The stability of the Sun's magnetic field
- D. The solar radio flux at Boulder, Colorado



G3A13 -- What does the A-index indicate?

- A. The relative position of sunspots on the surface of the Sun
- B. The amount of polarization of the Sun's electric field
- ➔ C. The long term stability of the Earth's geomagnetic field
- D. The solar radio flux at Boulder, Colorado



G3A14 -- How are radio communications usually affected by the charged particles that reach the Earth from solar coronal holes?

- A. HF communications are improved
- ➔ B. HF communications are disturbed
- C. VHF/UHF ducting is improved
- D. VHF/UHF ducting is disturbed



G3A15 -- How long does it take charged particles from coronal mass ejections to affect radio propagation on the Earth?

- A. 28 days
- B. 14 days
- C. 4 to 8 minutes
- ➔ D. 20 to 40 hours



G3A16 -- What is a possible benefit to radio communications resulting from periods of high geomagnetic activity?

- ➔ A. Auroras that can reflect VHF signals
- B. Higher signal strength for HF signals passing through the polar regions
- C. Improved HF long path propagation
- D. Reduced long delayed echoes



G3B03 -- Which of the following applies when selecting a frequency for lowest attenuation when transmitting on HF?

- ➔ A. Select a frequency just below the MUF
- B. Select a frequency just above the LUF
- C. Select a frequency just below the critical frequency
- D. Select a frequency just above the critical frequency



G3B04 -- What is a reliable way to determine if the MUF is high enough to support skip propagation between your station and a distant location on frequencies between 14 and 30 MHz?

- ➔ A. Listen for signals from an international beacon
- B. Send a series of dots on the band and listen for echoes from your signal
- C. Check the strength of TV signals from Western Europe
- D. Check the strength of signals in the MF AM broadcast band



G3B05 -- What usually happens to radio waves with frequencies below the MUF and above the LUF when they are sent into the ionosphere?

- ➔ A. They are bent back to the Earth
- B. They pass through the ionosphere
- C. They are amplified by interaction with the ionosphere
- D. They are bent and trapped in the ionosphere to circle the Earth



G3B06 -- What usually happens to radio waves with frequencies below the LUF?

- A. They are bent back to the Earth
- B. They pass through the ionosphere
- C. They are completely absorbed by the ionosphere
- D. They are bent and trapped in the ionosphere to circle the Earth



G3B07 -- What does LUF stand for?

- A. The Lowest Usable Frequency for communications between two points
- B. The Longest Universal Function for communications between two points
- C. The Lowest Usable Frequency during a 24 hour period
- D. The Longest Universal Function during a 24 hour period



G3B08 -- What does MUF stand for?

- A. The Minimum Usable Frequency for communications between two points
- ➔ B. The Maximum Usable Frequency for communications between two points
- C. The Minimum Usable Frequency during a 24 hour period
- D. The Maximum Usable Frequency during a 24 hour period



G3B11 -- What happens to HF propagation when the Lowest Usable Frequency (LUF) exceeds the Maximum Usable Frequency (MUF)?

- ➔ A. No HF radio frequency will support ordinary sky-wave communications over the path
- B. HF communications over the path are enhanced
- C. Double hop propagation along the path is more common
- D. Propagation over the path on all HF frequencies is enhanced



G3B12 -- What factor or factors affect the MUF?

- A. Path distance and location
- B. Time of day and season
- C. Solar radiation and ionospheric disturbances
- ➔ D. All of these choices are correct



Scatter Modes

- Scatter Characteristics
 - Localized areas in the ionosphere can reflect radio waves as well as refract them.
 - Direction of reflection is unpredictable.
 - Reflected signals are **MUCH** weaker than refracted signals & have a wavering sound.
 - Allows propagation above the MUF.



Scatter Modes

- Scatter Characteristics
 - Backscatter.
 - Signals can be reflected from uneven terrain at the far end of the path back towards the source.



Scatter Modes

- Near Vertical Incidence Sky-wave (NVIS)
 - At frequencies below the critical frequency, signals arriving at any angle are reflected.
 - Allows communications up to 200-300 hundred miles.
 - Eliminates the “skip zone”.



Scatter Modes

- Near Vertical Incidence Sky-wave (NVIS)
 - Select a frequency below the critical frequency but high enough that absorption in the D-region is not excessive.
 - Use a horizontally-polarized antenna mounted $1/8\lambda$ to $1/4\lambda$ above the ground.
 - 10' to 12' above the ground is best.
 - A grounded wire on the ground directly beneath the antenna improves signal by up to 6 dB.



G3C06 -- What is a characteristic of HF scatter signals?

- A. They have high intelligibility
- ➔ B. They have a wavering sound
- C. They have very large swings in signal strength
- D. All of these choices are correct



G3C07 -- What makes HF scatter signals often sound distorted?

- A. The ionospheric layer involved is unstable
- B. Ground waves are absorbing much of the signal
- C. The E-region is not present
- ➔ D. Energy is scattered into the skip zone through several different radio wave paths



G3C08 -- Why are HF scatter signals in the skip zone usually weak?

- ➔ A. Only a small part of the signal energy is scattered into the skip zone
- B. Signals are scattered from the magnetosphere which is not a good reflector
- C. Propagation is through ground waves which absorb most of the signal energy
- D. Propagation is through ducts in F region which absorb most of the energy



G3C09 -- What type of radio wave propagation allows a signal to be detected at a distance too far for ground wave propagation but too near for normal sky-wave propagation?

- A. Faraday rotation
- ➔ B. Scatter
- C. Sporadic-E skip
- D. Short-path skip



G3C10 -- Which of the following might be an indication that signals heard on the HF bands are being received via scatter propagation?

- A. The communication is during a sunspot maximum
- B. The communication is during a sudden ionospheric disturbance
- C. The signal is heard on a frequency below the Maximum Usable Frequency
- ➔ D. The signal is heard on a frequency above the Maximum Usable Frequency



G3C11 -- Which of the following antenna types will be most effective for skip communications on 40 meters during the day?

- A. Vertical antennas
- ➔ B. Horizontal dipoles placed between $1/8$ and $1/4$ wavelength above the ground
- C. Left-hand circularly polarized antennas
- D. Right-hand circularly polarized antenna



G3C13 -- What is Near Vertical Incidence Sky-wave (NVIS) propagation?

- A. Propagation near the MUF
- ➔ B. Short distance HF propagation using high elevation angles
- C. Long path HF propagation at sunrise and sunset
- D. Double hop propagation near the LUF



Questions?



Next Week

Chapter 9

Safety